

M.H
PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 1/20	A2	(11) International Publication Number: WO 00/01799 (43) International Publication Date: 13 January 2000 (13.01.00)
(21) International Application Number: PCT/DK99/00382 (22) International Filing Date: 2 July 1999 (02.07.99) (30) Priority Data: PA 1998 00878 3 July 1998 (03.07.98) DK 60/091,735 6 July 1998 (06.07.98) US (71) Applicant (for all designated States except US): CHR. HANSEN A/S [DK/DK]; Bøge Allé 10-12, DK-2970 Hørsholm (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): NILSSON, Dan [DK/DK]; Gefionsbakken 4A, DK-3060 Espergærde (DK). JANZEN, Thomas [DK/DK]; Ingemannsvej 30, 3. tv., DK-1964 Frederiksberg (DK). (74) Agent: PLOUGMANN, VINGTOFT & PARTNERS; Sankt Annæ Plads 11, P.O. Box 3007, DK-1021 Copenhagen K (DK).		(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHOD OF PREVENTING BACTERIOPHAGE INFECTION OF BACTERIAL CULTURES (57) Abstract Method of preventing bacteriophage infection of bacterial cultures comprising modified strains, wherein the cultures are completely resistant to bacteriophage attack and have retained their capability of being metabolically active. The method is useful in the manufacturing of food products, feed products or useful metabolite products.		